

Interacting open p-branes

Sumio Ishikawa*, Yasuhiro Iwama,
Tadashi Miyazaki and Motowo Yamanobe†
Department of Physics, Science University of Tokyo
Kagurazaka, Shinjuku-ku, Tokyo 162, Japan

December 1994

PACS 03.50.+z - Classical field theory

PACS 11.17.+y - Theories of strings and other extended objects

Abstract

The Kalb-Ramond action, derived for interacting strings through an action-at-a-distance force, is generalized to the case of interacting p -dimensional objects (p-branes) in D -dimensional space-time. The *open* p-brane version of the theory is especially taken up. On account of the existence of their boundary surface, the fields mediating interactions between open p-branes are obtained as *massive* gauge fields, quite in contrast to massless gauge ones for *closed* p-branes.

*E-mail: liszt@gradsun.ap.kagu.sut.ac.jp

†E-mail: yamanobe@gradsun.ap.kagu.sut.ac.jp

Some forty-five years ago, Feynman and Wheeler[1] presented an elegant idea that the action-at-a-distance (AD) force gives us a gauge field (photon) mediating electromagnetic interaction of a point particle. Use was made of this idea by Kalb and Ramond[2] to describe interacting strings[2][3]. They had the results that the fields mediating the mutual interactions of strings are either massless vector fields (for *closed* strings) or massive vector ones (for *open* strings).

The present authors generalized Kalb-Ramond's method to the case where extended objects of more than one-dimension are interacting with one another[4]. There *closed* p-branes were taken up and it was concluded that it is the *massless* gauge field that plays an essential role in the interaction of closed p-branes.

In the present paper, the authors will confront with *open* p-branes and construct the theory for the interacting open p-branes *à la* Kalb-Ramond's AD view. The existence of boundary surface in this case enforces us to take the equations on surface *ad hoc* into account. The difference here arises, so that one is to have a *massive* gauge field, as will soon be clear.

Now let us start with the following action for two p-branes, which is interacting each other through AD forces[4]

$$S = S_a^{\text{free}} + S_b^{\text{free}} + \int d^{p+1}\xi_a \int d^{p+1}\xi_b R_{ab}(X_a, X_b, X_{a,i}, X_{b,i}). \quad (1)$$

The p-brane is represented by the variables $X^\mu(\xi)$ ($\mu = 0, 1, \dots, D-1$) in the flat D -dimensional space-time. The world sheet of the p-brane is described by the variables ξ^i ($0 \leq \xi^i \leq l^i$; $i = 0, 1, \dots, p$). The subscript a or b labels each p-brane. S^{free} is the action for a free p-brane, and is taken to be the $(p+1)$ -dimensional world volume traced out by the p-brane

$$\begin{aligned} S^{\text{free}} &= - \int (-d\sigma \cdot d\sigma)^{\frac{1}{2}} \\ &= - \int d^{p+1}\xi (-\sigma \cdot \sigma)^{\frac{1}{2}}. \end{aligned} \quad (2)$$

The notations used in this paper are as follows:

$$d\sigma \cdot d\sigma \equiv d\sigma^{\mu_0\mu_1\cdots\mu_p} d\sigma_{\mu_0\mu_1\cdots\mu_p},$$

$$\begin{aligned}
\sigma \cdot \sigma &\equiv \sigma^{\mu_0 \mu_1 \dots \mu_p} \sigma_{\mu_0 \mu_1 \dots \mu_p}, \\
d\sigma^{\mu_0 \mu_1 \dots \mu_p} &\equiv d^{p+1} \xi \sigma^{\mu_0 \mu_1 \dots \mu_p}, \\
\sigma^{\mu_0 \mu_1 \dots \mu_p} &\equiv \frac{\partial(X^{\mu_0}, X^{\mu_1}, \dots, X^{\mu_p})}{\partial(\xi^0, \xi^1, \dots, \xi^p)}, \\
d^{p+1} \xi &\equiv d\xi^0 d\xi^1 \dots d\xi^p, \\
X_{,i}^\mu &\equiv \frac{\partial X^\mu}{\partial \xi^i}.
\end{aligned}$$

The R_{ab} satisfy

$$R_{ab} = R_{ba} \quad (3)$$

for symmetry requirements of the action. The variational principle for the variation $X_a^\mu(\xi) \rightarrow X_a^\mu(\xi) + \delta X_a^\mu(\xi)$ immediately yields the equations of motion

$$(p+1)D_a^{\mu_1 \mu_2 \dots \mu_p} \left[\frac{\sigma_{a\mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} \right] = \int d^{p+1} \xi_b \left[\left(\frac{\partial}{\partial X_a^{\mu_0}} - \sum_{i=0}^p \frac{\partial}{\partial \xi_a^i} \frac{\partial}{\partial X_{a,i}^{\mu_0}} \right) R_{ab} \right]. \quad (4)$$

Moreover, for our *open* p-brane, the boundary conditions on the surface add the following equations

$$\begin{aligned}
(p+1) \frac{\sigma_{a\mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} K_{ai}^{\mu_1 \mu_2 \dots \mu_p} &= - \int d^{p+1} \xi_b \frac{\partial R_{ab}}{\partial X_{a,i}^{\mu_0}}, \\
(\xi_a^i &= 0, l_a^i ; i = 1, 2, \dots, p).
\end{aligned} \quad (5)$$

Equation(5) constitutes one of the main characteristics of the open p-brane, and it will play an essential role later. Here and in what follows, we define K_i and D by

$$K_i^{\mu_1 \mu_2 \dots \mu_p} \equiv \frac{\partial \sigma^{\alpha \mu_1 \mu_2 \dots \mu_p}}{\partial X_{,i}^\alpha}, \quad (6)$$

$$D^{\mu_1 \mu_2 \dots \mu_p} \equiv \sum_{i=0}^p K_i^{\mu_1 \mu_2 \dots \mu_p} \frac{\partial}{\partial \xi^i}. \quad (7)$$

In the definition of K_i , we do not sum over the index α . We assume that R_{ab} depends on $X_{a,i}^\mu$ only through σ_a . Under this restriction, we rewrite Eq.(4) as

$$(p+1)D_a^{\mu_1 \mu_2 \dots \mu_p} \left[\frac{\sigma_{a\mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} \right] = \int d^{p+1} \xi_b \left[\left(\frac{\partial}{\partial X_a^{\mu_0}} - \frac{1}{p!} D_a^{\mu_1 \mu_2 \dots \mu_p} \frac{\partial}{\partial \sigma_a^{\mu_0 \mu_1 \dots \mu_p}} \right) R_{ab} \right], \quad (8)$$

and Eq.(5) as¹

$$\frac{\sigma_{a\mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} K_{ai}^{\mu_1 \mu_2 \dots \mu_p} = - \frac{1}{(p+1)!} \int d^{p+1} \xi_b K_{ai}^{\mu_1 \mu_2 \dots \mu_p} \frac{\partial R_{ab}}{\partial \sigma_a^{\mu_0 \mu_1 \dots \mu_p}}, \quad (9)$$

¹ Here care should be taken of the differentiation with respect to the antisymmetric tensor elements $\sigma^{\mu_0 \mu_1 \dots \mu_p}$. One is to differentiate in $\sigma^{\mu_0 \mu_1 \dots \mu_p}$, with every permutation of $\mu_1 \mu_2 \dots \mu_p$.

$$(\xi_a^i = 0, l_a^i).$$

Assume a concrete form of R_{ab} just in the same way as in the case of closed p-branes:

$$R_{ab} \equiv g_a g_b \sigma_a^{\mu_0 \mu_1 \dots \mu_p} \sigma_{b \mu_0 \mu_1 \dots \mu_p} G((X_a - X_b)^2), \quad (10)$$

with g_a and g_b , coupling constants, and G is some propagator in D -dimensional space-time.

Substitution of Eq.(10) into Eqs.(8) and (9) leads us to

$$(p+1)D_a^{\mu_1 \mu_2 \dots \mu_p} \left[\frac{\sigma_{a \mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} \right] = g_a \sigma_a^{\mu_1 \mu_2 \dots \mu_{p+1}} F_{b \mu_0 \mu_1 \dots \mu_{p+1}}(X_a), \quad (11)$$

and

$$\frac{\sigma_{a \mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} K_{ai}^{\mu_1 \mu_2 \dots \mu_p} = -g_a K_{ai}^{\mu_1 \mu_2 \dots \mu_p} \phi_{b \mu_0 \mu_1 \dots \mu_p}(X_a), \quad (12)$$

$$(\xi_a^i = 0, l_a^i),$$

where F_b and ϕ_b are antisymmetric tensors defined by

$$F_b^{\mu_0 \mu_1 \dots \mu_{p+1}}(X) \equiv \sum_{i=0}^{p+1} (-1)^{(p+1) \cdot i} \frac{\partial}{\partial X_{\mu_i}} \phi_b^{\mu_{i+1} \mu_{i+2} \dots \mu_{p+1} \mu_0 \mu_1 \dots \mu_{i-1}}(X), \quad (13)$$

$$\phi_b^{\mu_0 \mu_1 \dots \mu_p}(X) \equiv g_b \int d^{p+1} \xi_b \sigma_b^{\mu_0 \mu_1 \dots \mu_p} G((X - X_b)^2). \quad (14)$$

Now, let us investigate the conservation of a current for field ϕ_b and the gauge invariance of the action.

First, we consider the conservation of a current. In general, G obeys, being a propagator,

$$\left(\frac{\partial^2}{\partial X_\mu \partial X^\mu} + m^2 \right) G(X^2) = -C \delta^{(D)}(X), \quad (15)$$

where C is a dimensionless constant and m is to be identified with the mass of the field. By Eq.(15), we obtain

$$\left(\frac{\partial^2}{\partial X_\mu \partial X^\mu} + m^2 \right) \phi_b^{\mu_0 \mu_1 \dots \mu_p}(X) = -C j_b^{\mu_0 \mu_1 \dots \mu_p}(X), \quad (16)$$

with

$$j_b^{\mu_0 \mu_1 \dots \mu_p}(X) \equiv g_b \int d^{p+1} \xi_b \sigma_b^{\mu_0 \mu_1 \dots \mu_p} \delta^{(D)}(X - X_b). \quad (17)$$

We further find, with a bit of algebra, that

$$\frac{\partial}{\partial X^{\mu_0}} \phi_b^{\mu_0 \mu_1 \dots \mu_p}(X) = -g_b \sum_{i=0}^p \int d\xi_b^0 d\xi_b^1 \dots d\hat{\xi}_b^i \dots d\xi_b^p \left[K_{bi}^{\mu_1 \mu_2 \dots \mu_p} G((X - X_b)^2) \right]_{\xi_b^i=0}^{\xi_b^i=l_b^i}. \quad (18)$$

Here $d\hat{\xi}^i$ means exclusion of $d\xi^i$. On account of the existence of boundary surface, the divergence of ϕ_b does not vanish. From Eqs.(16) and (18) follows the equation

$$\frac{\partial}{\partial X^{\mu_0}} j_b^{\mu_0 \mu_1 \dots \mu_p}(X) = -g_b \sum_{i=0}^p \int d\xi_b^0 d\xi_b^1 \dots d\hat{\xi}_b^i \dots d\xi_b^p \left[K_{bi}^{\mu_1 \mu_2 \dots \mu_p} \delta^{(D)}(X - X_b) \right]_{\xi_b^i=0}^{\xi_b^i=l_b^i}, \quad (19)$$

which shows that the current j_b would be conserved if the p-brane was closed. The modification of the current for the open p-brane will be done after the introduction of a gauge transformation.

Secondly, we examine the gauge invariance of the action, defining a gauge transformation by

$$\phi^{\mu_0 \mu_1 \dots \mu_p}(X) \rightarrow \phi^{\mu_0 \mu_1 \dots \mu_p}(X) + \sum_{i=0}^p (-1)^{p \cdot i} \frac{\partial}{\partial X^{\mu_i}} \Lambda^{\mu_{i+1} \mu_{i+2} \dots \mu_p \mu_0 \mu_1 \dots \mu_{i-1}}, \quad (20)$$

with an arbitrary antisymmetric tensor Λ . Instead of the whole action S , we take up

$$\begin{aligned} S^{\text{int}} &\equiv \int d^{p+1} \xi_a \int d^{p+1} \xi_b R_{ab} \\ &= g_a \int d^{p+1} \xi_a \sigma_a^{\mu_0 \mu_1 \dots \mu_p} \phi_{b \mu_0 \mu_1 \dots \mu_p}, \end{aligned} \quad (21)$$

i.e., the ϕ -depending part S^{int} of S . Under the transformation (20), S^{int} transforms as follows:

$$S^{\text{int}} \rightarrow S^{\text{int}} + (p+1) g_a \int d^{p+1} \xi_a D_a^{\mu_1 \mu_2 \dots \mu_p} \Lambda_{\mu_1 \mu_2 \dots \mu_p} \quad (22)$$

$$= S^{\text{int}} + (p+1) g_a \int d\xi_a^0 d\xi_a^1 \dots d\hat{\xi}_a^i \dots d\xi_a^p \left[\sum_{i=0}^p K_{ai}^{\mu_1 \mu_2 \dots \mu_p} \Lambda_{\mu_1 \mu_2 \dots \mu_p} \right]_{\xi_a^i=0}^{\xi_a^i=l_a^i}. \quad (23)$$

The second term of the right-hand side in Eq.(23) does not vanish, here also, due to the existence of boundary surface. Note that it vanishes for *closed* p-branes. With Eq.(23), one immediately sees that S^{int} is not invariant under the transformation (20). Although the equations of motion (11) are invariant under this gauge transformation, the equations of boundary surface (12) are not. The latter become

$$\frac{\sigma_{a \mu_0 \mu_1 \dots \mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} K_{ai}^{\mu_1 \mu_2 \dots \mu_p} = -g_a K_{ai}^{\mu_1 \mu_2 \dots \mu_p} \left[\phi_{b \mu_0 \mu_1 \dots \mu_p}(X_a) + \sum_{i=0}^p (-1)^{p \cdot i} \frac{\partial}{\partial X^{\mu_i}} \Lambda_{\mu_{i+1} \mu_{i+2} \dots \mu_p \mu_0 \mu_1 \dots \mu_{i-1}} \right], \quad (24)$$

$$(\xi_a^i = 0, l_a^i).$$

To recover the invariance of the action, we therefore add another term to the action S . Based on Eqs.(23) and (24), we take up as an additional action

$$S^{\text{add}} \equiv -(p+1)e_a e_b \int d^{p+1}\xi_a \int d^{p+1}\xi_b D_a^{\mu_1\mu_2\cdots\mu_p} D_{b\mu_1\mu_2\cdots\mu_p} G((Xa - Xb)^2). \quad (25)$$

It is straightforward, under the variation in X_a^μ , that S^{add} does not change the equations of motion. Furthermore, Eq.(25) is rewritten as follows:

$$S^{\text{add}} = (p+1)e_a \int d^{p+1}\xi_a D_a^{\mu_1\mu_2\cdots\mu_p} B_{b\mu_1\mu_2\cdots\mu_p}, \quad (26)$$

with

$$B_b^{\mu_1\mu_2\cdots\mu_p}(X) \equiv -e_b \sum_{i=0}^p \int d\xi_b^0 d\xi_b^1 \cdots d\hat{\xi}_b^i \cdots d\xi_b^p \left[K_{bi}^{\mu_1\mu_2\cdots\mu_p} G((X - X_b)^2) \right]_{\xi_b^i=0}^{\xi_b^i=l_b^i}. \quad (27)$$

Equations (26) and (27) trivially teach us that S^{add} vanishes when p-branes a and b are closed.

In view of Eqs.(22) and (26), we define a new gauge transformation

$$\begin{aligned} \phi^{\mu_0\mu_1\cdots\mu_p}(X) &\rightarrow \phi^{\mu_0\mu_1\cdots\mu_p}(X) + \sum_{i=0}^p (-1)^{p \cdot i} \frac{\partial}{\partial X_{\mu_i}} \Lambda^{\mu_{i+1}\mu_{i+2}\cdots\mu_p\mu_0\mu_1\cdots\mu_{i-1}}, \\ B^{\mu_1\mu_2\cdots\mu_p}(X) &\rightarrow B^{\mu_1\mu_2\cdots\mu_p} - \frac{g}{e} \Lambda^{\mu_1\mu_2\cdots\mu_p}. \end{aligned} \quad (28)$$

It is easy to see that these transformations make the action $S + S^{\text{add}}$ invariant.

Under the variation in X_a^μ , the action $S + S^{\text{add}}$ yields the following equations of boundary surface

$$\frac{\sigma_{a\mu_0\mu_1\cdots\mu_p}}{(-\sigma_a \cdot \sigma_a)^{\frac{1}{2}}} K_{ai}^{\mu_1\mu_2\cdots\mu_p} = - \left[g_a \phi_{b\mu_0\mu_1\cdots\mu_p}(X_a) + e_a \sum_{i=0}^p (-1)^{p \cdot i} \frac{\partial}{\partial X_a^{\mu_i}} B_{b\mu_{i+1}\mu_{i+2}\cdots\mu_p\mu_0\mu_1\cdots\mu_{i-1}} \right] K_{ai}^{\mu_1\mu_2\cdots\mu_p}, \quad (29)$$

$$(\xi_a^i = 0, l_a^i).$$

These equations of boundary surface are invariant under the transformations (28).

The Lagrangian density for the fields ϕ and B is given by

$$\mathcal{L} = -\frac{1}{2(p+2)!} F^{\mu_0\mu_1\cdots\mu_{p+1}} F_{\mu_0\mu_1\cdots\mu_{p+1}}$$

$$\begin{aligned}
& + \frac{1}{2(p+1)!} \left[\sum_{i=0}^p (-1)^{p-i} \frac{\partial}{\partial X^{\mu_i}} B_{\mu_{i+1}\mu_{i+2}\dots\mu_p\mu_0\mu_1\dots\mu_{i-1}} \right]^2 \\
& + \frac{1}{2(p+1)!} \frac{g^2}{e^2} \phi^{\mu_0\mu_1\dots\mu_p} \phi_{\mu_0\mu_1\dots\mu_p} \\
& + \frac{1}{(p+1)!} \frac{g}{e} \phi^{\mu_0\mu_1\dots\mu_p} \left[\sum_{i=0}^p (-1)^{p-i} \frac{\partial}{\partial X^{\mu_i}} B_{\mu_{i+1}\mu_{i+2}\dots\mu_p\mu_0\mu_1\dots\mu_{i-1}} \right], \tag{30}
\end{aligned}$$

which is also invariant under the transformations (28). The variational principle for this Lagrangian $\mathcal{L}$ with respect to ϕ and B leads to the equation of motion for B :

$$\frac{\partial^2}{\partial X_{\mu_0} \partial X^{\mu_0}} B^{\mu_1\mu_2\dots\mu_p} = - \sum_{i=1}^p (-1)^{p-i} \frac{\partial}{\partial X_{\mu_i}} \frac{\partial}{\partial X^{\mu_0}} B^{\mu_{i+1}\mu_{i+2}\dots\mu_p\mu_0\mu_1\dots\mu_{i-1}} - \frac{g}{e} \frac{\partial}{\partial X^{\mu_0}} \phi^{\mu_0\mu_1\dots\mu_p}, \tag{31}$$

and that for ϕ :

$$\begin{aligned}
\frac{\partial^2}{\partial X_{\mu_0} \partial X^{\mu_0}} \phi^{\mu_1\mu_2\dots\mu_{p+1}} & = - \sum_{i=1}^{p+1} (-1)^{(p+1)-i} \frac{\partial}{\partial X_{\mu_i}} \frac{\partial}{\partial X^{\mu_0}} \phi^{\mu_{i+1}\mu_{i+2}\dots\mu_{p+1}\mu_0\mu_1\dots\mu_{i-1}} \\
& - \frac{g}{e} \sum_{i=1}^{p+1} (-1)^{p-(i+1)} \frac{\partial}{\partial X_{\mu_i}} B^{\mu_{i+1}\mu_{i+2}\dots\mu_{p+1}\mu_1\mu_2\dots\mu_{i-1}} \\
& - \frac{g^2}{e^2} \phi^{\mu_1\mu_2\dots\mu_{p+1}}. \tag{32}
\end{aligned}$$

We next investigate the conservation of a current for B_b . Comparing Eq.(27) with Eq.(18), we immediately find that ϕ_b and B_b satisfy a gauge condition

$$\frac{\partial}{\partial X^{\mu_0}} \phi_b^{\mu_0\mu_1\dots\mu_p}(X) = \frac{g_b}{e_b} B_b^{\mu_1\mu_2\dots\mu_p}(X). \tag{33}$$

Furthermore, B_b obeys

$$\frac{\partial}{\partial X^{\mu_1}} B_b^{\mu_1\mu_2\dots\mu_p}(X) = 0. \tag{34}$$

Therefore, we define a new current J_b for B_b by

$$J_b^{\mu_1\mu_2\dots\mu_p} \equiv e_b \sum_{i=1}^p \int d\xi_b^0 d\xi_b^1 \dots d\hat{\xi}_b^i \dots d\xi_b^p \left[K_{bi}^{\mu_1\mu_2\dots\mu_p} \delta^{(D)}(X - X_b) \right]_{\xi_b^i=0}^{\xi_b^i=l_b^i}, \tag{35}$$

by which we have

$$\left(\frac{\partial^2}{\partial X_{\mu} \partial X^{\mu}} + m^2 \right) B_b^{\mu_1\mu_2\dots\mu_p}(X) = -C J_b^{\mu_1\mu_2\dots\mu_p}(X). \tag{36}$$

According to the vanishing of the divergence of B_b , J_b satisfies the conservation law of a current

$$\frac{\partial}{\partial X^{\mu_1}} J_b^{\mu_1\mu_2\dots\mu_p} = 0. \tag{37}$$

It follows from Eqs.(16), (36) and (33) that

$$J_b^{\mu_1\mu_2\cdots\mu_p}(X) = \frac{e_b}{g_b} \frac{\partial}{\partial X^{\mu_0}} j_b^{\mu_0\mu_1\cdots\mu_p}(X). \quad (38)$$

This shows that the new conserved current J_b instead of the nonconserved one j_b is *physical*. Note that the divergence of j_b enters into Eq.(38).

With gauge conditions (33) and (34), Eq.(31) becomes

$$\frac{\partial^2}{\partial X_{\mu_0} \partial X^{\mu_0}} B^{\mu_1\mu_2\cdots\mu_p} = - \left(\frac{g}{e}\right)^2 B^{\mu_1\mu_2\cdots\mu_p}, \quad (39)$$

which means B is a field with mass

$$m = \frac{g}{e}. \quad (40)$$

Similarly, the equation of motion for ϕ is

$$\frac{\partial^2}{\partial X_{\mu_0} \partial X^{\mu_0}} \phi^{\mu_1\mu_2\cdots\mu_{p+1}} = - \left(\frac{g}{e}\right)^2 \phi^{\mu_1\mu_2\cdots\mu_{p+1}}, \quad (41)$$

and ϕ is considered as a field with mass $m = g/e$.

Now we combine ϕ with B to introduce a new field

$$\psi^{\mu_0\mu_1\cdots\mu_p}(X) \equiv \phi^{\mu_0\mu_1\cdots\mu_p}(X) + \frac{e}{g} \sum_{i=0}^p (-1)^{p-i} \frac{\partial}{\partial X^{\mu_i}} B^{\mu_{i+1}\mu_{i+2}\cdots\mu_p\mu_0\mu_1\cdots\mu_{i-1}}(X). \quad (42)$$

It is clear that this field ψ is invariant under the transformations (28). In terms of ψ , we can rewrite $\mathcal{L}$ as

$$\mathcal{L} = -\frac{1}{2(p+2)!} F^{\mu_0\mu_1\cdots\mu_{p+1}} F_{\mu_0\mu_1\cdots\mu_{p+1}} + \frac{1}{2(p+1)!} \frac{g^2}{e^2} \psi^{\mu_0\mu_1\cdots\mu_p} \psi_{\mu_0\mu_1\cdots\mu_p}. \quad (43)$$

The gauge invariance of ψ also leads to the invariance of $\mathcal{L}$.

The gauge conditions (18) and (34) induce, for ψ ,

$$\frac{\partial}{\partial X^{\mu_0}} \psi^{\mu_0\mu_1\cdots\mu_p} = 0. \quad (44)$$

And the equation of motion for ψ is obtained as

$$\frac{\partial^2}{\partial X_{\mu_0} \partial X^{\mu_0}} \psi^{\mu_1\mu_2\cdots\mu_{p+1}} = - \left(\frac{g}{e}\right)^2 \psi^{\mu_1\mu_2\cdots\mu_{p+1}}. \quad (45)$$

It shows that ψ is again a field with mass $m = g/e$.

Thus we have obtained a massive gauge-invariant field ψ . The total action, which describes the open p-brane and its interaction with the field ψ , is given by

$$S^{\text{total}} = - \int d^{p+1} \xi (-\sigma \cdot \sigma)^{\frac{1}{2}} + g \int d^{p+1} \xi \sigma^{\mu_0 \mu_1 \dots \mu_p} \psi_{\mu_0 \mu_1 \dots \mu_p} + \int d^D X \mathcal{L}. \quad (46)$$

Now we come to our conclusions. As we have discussed above, it is the existence of the boundary surface that makes a difference between closed and open p-branes. Closed p-branes are described by the equations of motion (11) without equations of boundary surface. Consequently, their interactions are regarded as mediated by the massless field ϕ defined by Eq.(14)[4]. But when ϕ is introduced in case of open p-branes, two difficulties occur owing to this existence of the boundary surface of world volumes. The one is that the current j , defined by Eq.(17), for ϕ is not to be conserved, the other is that the action S is not gauge invariant under the transformation (20). To overcome these difficulties we have added the new term S^{add} to the action S , and defined the new gauge transformations by Eq.(28). The introduction of them brings us to that of a new field B . This further enforces us to have the action $S + S^{\text{add}}$ instead of S , by which the action has been recovered its gauge invariance. We find that the current J for the field B , which is regarded as the divergence of the current j , is conserved with Eqs.(37) and (38).

As a consequence, we have obtained the field ψ , which is constructed by the combination of the fields ϕ and B , as a medium of interaction of the open p-branes. In contrast to the case of the closed p-branes the field ψ has clearly a mass $m = g/e$; besides, it is gauge invariant under the transformation (28).

One of the authors (M.Y.) would like to thank Iwanami Fûjukai for financial support.

References

- [1] R. P. Feynman and J. A. Wheeler, Rev. Mod. Phys. **24**, 425 (1949).
- [2] M. Kalb and P. Ramond, Phys. Rev. **D 9**, 2273 (1974).
- [3] P. Ramond, Phys. Rev. **D 7**, 449 (1973).
- [4] M. Yamanobe, S. Ishikawa, Y. Iwama, T. Miyazaki, K. Yamamoto and R. Yoshida, preprint of Science University of Tokyo (1994), hep-th/9408122.